

Checklist for Prevention of Central Line Associated Blood Stream Infections

Based on 2011 CDC guideline for prevention of intravascular catheter-associated bloodstream infections:
<http://www.cdc.gov/hicpac/pdf/guidelines/bsi-guidelines-2011.pdf>

For Clinicians:

Promptly remove unnecessary central lines

- Perform daily audits to assess whether each central line is still needed

Follow proper insertion practices

- Perform hand hygiene before insertion
- Adhere to aseptic technique
- Use maximal sterile barrier precautions (i.e., mask, cap, gown, sterile gloves, and sterile full-body drape)
- Perform skin antisepsis with >0.5% chlorhexidine with alcohol
- Choose the best site to minimize infections and mechanical complications
 - Avoid femoral site in adult patients
- Cover the site with sterile gauze or sterile, transparent, semipermeable dressings

Handle and maintain central lines appropriately

- Comply with hand hygiene requirements
- Scrub the access port or hub immediately prior to each use with an appropriate antiseptic (e.g., chlorhexidine, povidone iodine, an iodophor, or 70% alcohol)
- Access catheters only with sterile devices
- Replace dressings that are wet, soiled, or dislodged
- Perform dressing changes under aseptic technique using clean or sterile gloves

For Facilities:

- Empower staff to stop non-emergent insertion if proper procedures are not followed
- "Bundle" supplies (e.g., in a kit) to ensure items are readily available for use
- Provide the checklist above to clinicians, to ensure all insertion practices are followed
- Ensure efficient access to hand hygiene
- Monitor and provide prompt feedback for adherence to hand hygiene
<http://www.cdc.gov/handhygiene/Measurement.html>
- Provide recurring education sessions on central line insertion, handling and maintenance

Supplemental strategies for consideration:

- 2% Chlorhexidine bathing
- Antimicrobial/Antiseptic-impregnated catheters
- Chlorhexidine-impregnated dressings